arXiv:1302.4190v1 [math.GN] 18 Feb 2013

ON PARATOPOLOGICAL GROUPS

FUCAI LIN AND CHUAN LIU*

Abstract. In this paper, we firstly construct a Hausdorff non-submetrizable paratopological group G in which every point is a G_δ-set, which gives a negative answer to Arhangel'skiĭ and Tkachenko's question [Topological Groups and Related Structures, Atlantis Press and World Sci., 2008]. We prove that each first-countable Abelian paratopological group is submetrizable. Moreover, we discuss developable paratopological groups and construct a non-metrizable, Moore paratopological group. Further, we prove that a regular, countable, locally k_ω-paratopological group is a discrete topological group or contains a closed copy of S_ω. Finally, we discuss some properties on non-H-closed paratopological groups, and show that Sorgenfrey line is not H-closed, which gives a negative answer to Arhangel'skiĭ and Tkachenko's question [Topological Groups and Related Structures, Atlantis Press and World Sci., 2008]. Some questions are posed.

1. Introduction

A *semitopological group* G is a group G with a topology such that the product map of $G\times G$ into G is separately continuous. A *paratopological group* G is a group G with a topology such that the product map of $G\times G$ into G is jointly continuous. If G is a paratopological group and the inverse operation of G is continuous, then G is called a *topological group.* However, there exists a paratopological group which is not a topological group; Sorgenfrey line ([16, Example 1.2.2]) is such an example. Paratopological groups were discussed and many results have been obtained [7, 8, 6, 14, 22, 23, 24, 19].

Proposition 1.1. [29] For a group with topology (G,τ) the following conditions are equivalent:

1. G is a paratopological group;
2. The following Pontrjagin conditions for basis $\mathscr{B}=\mathscr{B}_\tau$ of the neutral element e of G are satisfied.
   - (a) $(\forall U,V\in\mathscr{B})(\exists W\in\mathscr{B})$: $W\subset U\cap V$;
   - (b) $(\forall U\in\mathscr{B})(\exists V\in\mathscr{B})$: $V^2\subset U$;
   - (c) $(\forall U\in\mathscr{B})(\forall x\in U)(\exists V\in\mathscr{B})$: $Vx\subset U$;
   - (d) $(\forall U\in\mathscr{B})(\forall x\in G)(\exists V\in\mathscr{B})$: $xVx^{-1}\subset U$;

2000 *Mathematics Subject Classification.* 54E20; 54E35; 54H11; 22A05.

Key words and phrases. Paratopological groups; submetrizable; ω-narrow; Abelian groups; developable; quasi-developable; k_ω-paratopological groups; H-closed paratopological groups; pseudocompact.

The first author is supported by the NSFC (No. 10971185, 10971186) and the Natural Science Foundation of Fujian Province (No. 2011J05013) of China.

*corresponding author.

The paratopological group G is Hausdorff if and only if
(e) $\bigcap\{UU^{-1} : U \in \mathscr{B}\} = \{e\}$;

The paratopological group G is a topological group if and only if
(f) $(\forall U \in \mathscr{B})(\exists V \in \mathscr{B})$: $V^{-1} \subset U$.

In this paper, we mainly discuss the following questions.

Question 1.2. [8, Open problem 3.3.1] *Suppose that G is a Hausdorff (regular) paratopological group in which every point is a G_δ-set. Is G submetrizable?*

Question 1.3. [8, Open problem 5.7.2] *Let G be a regular first-countable ω-narrow paratopological group . Is G submetrizable?*

Question 1.4. [5, Problem 20] *Is every regular first countable (Abelian) paratopological group submetrizable?*

Question 1.5. [5, Problem 22] *Is it true that every regular first countable (Abelian) paratopological group G has a zero-set diagonal*[1]*?*

Question 1.6. [5, Problem 21] *Is every regular first countable (Abelian) paratopological group Dieudonné complete?*

Question 1.7. [8, Open problem 3.4.3] *Let G be a regular ω-narrow first-countable paratopological group. Does there exist a continuous isomorphism of G onto a regular (Hausdorff) second-countable paratopological group?*

Question 1.8. [24] *Is a regular symmetrizable paratopological group metrizable?*

Question 1.9. [8, Open problem 5.7.5] *Is every paratopological group, which is Moore space, metrizable?*

Question 1.10. [8, Open problem 3.6.5] *Must the Sorgenfrey line S be closed in every Hausdorff paratopological group containing it as a paratopological subgroup?*

We shall give negative answers to Questions 1.2, 1.8, 1.10, and 4.2, and give a partial answer to Question 1.3. Moreover, we shall also give affirmative answers to Questions 1.4, 1.5, 1.7 and 1.6 when the group G is Abelian.

2. Preliminaries

Definition 2.1. Let $\mathscr{P} = \bigcup_{x\in X} \mathscr{P}_x$ be a cover of a space X such that for each $x \in X$, (a) if $U, V \in \mathscr{P}_x$, then $W \subset U \cap V$ for some $W \in \mathscr{P}_x$; (b) the family $\mathscr{P}_x$ is a network of x in X, i.e., $x \in \bigcap \mathscr{P}_x$, and if $x \in U$ with U open in X, then $P \subset U$ for some $P \in \mathscr{P}_x$.

The family $\mathscr{P}$ is called a *weak base* for X [1] if, for every $A \subset X$, the set A is open in X whenever for each $x \in A$ there exists $P \in \mathscr{P}_x$ such that $P \subset A$. The space X is *weakly first-countable* if $\mathscr{P}_x$ is countable for each $x \in X$.

Definition 2.2. (1) A space X is called an S_ω-space if X is obtained by identifying all the limit points from a topological sum of countably many convergent sequences;

---

[1] We say that a space X has a *zero-set diagonal* if the diagonal in $X \times X$ is a zero-set of some continuous real-valued function on $X \times X$.

(2) A space X is called an S_2-space (*Arens' space*) if $X = \{\infty\} \cup \{x_n : n \in \mathbb{N}\} \cup \{x_n(m) : m, n \in \mathbb{N}\}$ and the topology is defined as follows: Each $x_n(m)$ is isolated; a basic neighborhood of x_n is $\{x_n\} \cup \{x_n(m) : m > k,$ for some $k \in \mathbb{N}\}$; a basic neighborhood of ∞ is $\{\infty\} \cup (\bigcup\{V_n : n > k \text{ for some } k \in \mathbb{N}\})$, where V_n is a neighborhood of x_n.

Definition 2.3. Let X be a space and $\{\mathscr{P}_n\}_n$ a sequence of collections of open subsets of X.

1. $\{\mathscr{P}_n\}_n$ is called a *quasi-development* [10] for X if for every $x \in U$ with U open in X, there exists an $n \in \mathbb{N}$ such that $x \in \mathrm{st}(x, \mathscr{P}_n) \subset U$.
2. $\{\mathscr{P}_n\}_n$ is called a *development* [30] for X if $\{\mathrm{st}(x, \mathscr{P}_n)\}_n$ is a neighborhood base at x in X for each point $x \in X$.
3. X is called *quasi-developable* (resp. *developable*), if X has a quasi-development (resp. *development*).
4. X is called *Moore*, if X is regular and developable.

A subset B of a paratopological group G is called *ω-narrow* in G if, for each neighborhood U of the neutral element of G, there is a countable subset F of G such that $B \subset FU \cap UF$.

A space X is called a *submetrizable* space if it can be mapped onto a metric space by a continuous one-to-one map. A space X is called a *subquasimetrizable* space if it can be mapped onto a quasimetric space by a one-to-one map.

All spaces are T_0 unless stated otherwise. The notations $\mathbb{R}, \mathbb{Q}, \mathbb{P}, \mathbb{N}, \mathbb{Z}$ are real numbers, rational numbers, irrational numbers, natural numbers and integers respectively. The letter e denotes the neutral element of a group. Readers may refer to [8, 16, 17] for notations and terminology not explicitly given here.

3. Submetrizablity of first-countable paratopological groups

In this section, we firstly give a negative answer to Question 1.2, then a answer to Question 1.3. We also give affirmative answers to Questions 1.4, 1.5, 1.7 and 1.6 when G is Abelian.

Proposition 3.1. [27] The following conditions are equivalent for an arbitrary space X.

1. The space X is submetrizable.
2. The free paratopological group $F_p(X)$ is submetrizable.
3. The free Abelian paratopological group $A_p(X)$ is submetrizable.

Proposition 3.2. [27] The following conditions are equivalent for an arbitrary space X.

1. The space X is subquasimetrizable.
2. The free paratopological group $F_p(X)$ is subquasimetrizable.
3. The free Abelian paratopological group $A_p(X)$ is subquasimetrizable.

Example 3.3. There exist a Hausdorff paratopological group G in which every point is a G_δ-set, and G is not submetrizable.

Proof. Let X be the lexicographically ordered set $X = (\mathbb{R} \times \{0\}) \cup (\mathbb{P} \times \mathbb{Z})$. Then X is a non-metrizable linearly ordered topological space without G_δ-diagonal ([11, Example 2.4]), hence X is not submetrizable. However, X is quasi-developable [25]. It is well known that quasi-developability in a generalized ordered space is equivalent to the existence of a σ-disjoint base, or of a σ-point-finite base [12, Theorem 4.2]. Hence X has σ-point finite base. Therefore, X is quasi-metrizable since a space with a σ-point finite base is quasi-metrizable [17, Page 489]. Let G be the free Abelian paratopological group $A_p(X)$ over X. Since a totally order space endowed the order topology is Tychonoff, then X is Tychonoff. It follows from [27, Proposition 3.8] that G is Hausdorff. By Propositions 3.1 and 3.2, G is subquasimetrizable and non-submetrizable. Since G is subquasimetrizable, every singleton of G is a G_δ-set. $\square$

Next we partially answer Question 1.3.

The *weak extent* [9] of a space X, denoted by $\omega e(X)$, is the least cardinal number κ such that for every open cover $\mathscr{U}$ of X there is a subset A of X of cardinality no greater than κ such that $\mathrm{st}(A;\mathscr{U}) = X$, where $\mathrm{st}(A;\mathscr{U}) = \bigcup\{U : U \in \mathscr{U}, U \cap A \neq \emptyset\}$. If X is separable, then $we(X) = \omega$.

Theorem 3.4. [9] *If X^2 has countable weak extent and a regular G_δ-diagonal*[2]*, then X condenses onto a second countable Hausdorff space.*

Theorem 3.5. [19] *Each ω-narrow first-countable paratopological group is separable.*

Theorem 3.6. *If G is a regular ω-narrow first-countable paratopological group, then G condenses onto a second countable Hausdorff space..*

Proof. It is straightforward to prove that the product of two ω-narrow paratopological groups is an ω-narrow paratopological group. Then G^2 is an ω-narrow first-countable paratopological group, and hence G^2 is separable by Theorem 3.5. Then G^2 has countable weak extent. Moreover, it follows from [22] that G has a regular G_δ-diagonal. Therefore, G condenses onto a second countable Hausdorff space by Theorem 3.4. $\square$

Corollary 3.7. *Let (G,τ) be a regular ω-narrow first-countable paratopological group. There exists a continuous isomorphism of G onto a Hausdorff second-countable space.*

A paratopological group G with a base at the neutral element $\mathscr{B}$ is a *SIN-group* (Small Invariant Neighborhoods), if for each $U \in \mathscr{B}$ there exists a $V \in \mathscr{B}$ such that $xVx^{-1} \subset U$ for each $x \in G$.

Theorem 3.8. *If (G,τ) is a Hausdorff SIN first-countable paratopological group, then G is submetrizable.*

Proof. Let $\{U_n : n \in \mathbb{N}\}$ be a countable local base of (G,τ) at the neutral element e, where $U_{n+1} \subset U_n$ for each $n \in \mathbb{N}$.

For $x \in G$, let $\mathcal{B}_x = \{xU_nU_n^{-1} : n \in \mathbb{N}\}$. Then $\{\mathcal{B}_x\}_{x\in G}$ has the following properties.

(BP1) For every $x \in G$, $\mathcal{B}_x \neq \emptyset$ and for every $U \in \mathcal{B}_x, x \in U$.

(BP2) If $x \in U \in \mathcal{B}_y$, then there exists a $V \in \mathcal{B}_x$ such that $V \subset U$

In fact, if $x \in U = yU_iU_i^{-1} \in \mathcal{B}_y$, then $x = yu_1u_2^{-1}$ for some $u_1, u_2 \in U_i$. Pick $U_j, U_k \in \{U_n : n \in \mathbb{N}\}$ such that $U_k \subset U_i$, $u_1U_k \subset U_i, U_ju_2 \subset U_k, u_2^{-1}U_ju_2 \subset U_k$.

---

[2] A space X is said to have a *regular G_δ-diagonal* if the diagonal $\Delta = \{(x,x) : x \in X\}$ can be represented as the intersection of the closures of a countable family of open neighborhoods of Δ in $X \times X$.

Then $xU_jU_j^{-1} = yu_1u_2^{-1}U_jU_j^{-1} \subset yu_1U_ku_2^{-1}U_j^{-1} \subset yU_i(U_ju_2)^{-1} \subset yU_iU_k^{-1} \subset yU_iU_i^{-1} = U$.

(BP3) For any $V_1, V_2 \in \mathcal{B}_x$ there exists a $V \in \mathcal{B}_x$ such that $V \subset V_1 \cap V_2$.

Let τ^* be the topology generated by the neighborhood system $\{\mathcal{B}_x\}_{x\in G}$. Obviously, the topology of (G, τ^*) is coarser than (G, τ) and it is first-countable. We prove that (G, τ^*) is a Hausdorff topological group.

It is easy to see that (a), (d) and (f) in Proposition 1.1 are satisfied. (BP2) implies (c). We check conditions (b) and (d).

Fix $n \in \mathbb{N}$. Then there is $k > n$ such that $U_k^2 \subset U_n$ since (G, τ) is a paratopological group. G is a SIN-group, there exists an $m > k$ such that $xU_mx^{-1} \subset U_k$ for each $x \in G$, and hence $U_m^{-1}U_m \subset U_kU_m^{-1}$. Then $U_mU_m^{-1}U_mU_m^{-1} \subset U_mU_kU_m^{-1}U_k^{-1} \subset U_kU_kU_k^{-1}U_k^{-1} \subset U_nU_n^{-1}$. Hence (b) is satisfied.

Since $\{U_n : n \in \mathbb{N}\}$ is a network of G at e, $\bigcap_{n\in\mathbb{N}} U_nU_n^{-1} = \{e\}$ by [2, Proposition 3.4, 3.5]. Therefore, (G, τ^*) is Hausdorff, and hence it is Tychonoff.

Since first-countable topological group metrizable, we have (G, τ^*) is metrizable. Therefore, G is submetrizable. □

It is well known that all submetrizable spaces have a zero-set diagonal. Therefore, Theorem 3.8 gives a partial answer to Question 1.5.

Corollary 3.9. *If (G, τ) is a Hausdorff Abelian first-countable paratopological group, then G is submetrizable.*

Indeed, we have the following more stronger result.

Theorem 3.10. *If (G, τ) is a Hausdorff Abelian paratopological group with a countable π-character, then G is submetrizable.*

Proof. Let $\mathscr{B} = \{U_\alpha : \alpha < \kappa\}$ be a local base at the neutral element e. It follows from the proof of Theorem 3.8 that the family $\{U_\alpha U_\alpha^{-1} : \alpha < \kappa\}$ is a local base at point e in the Tychonoff topological group (G, τ^*).

Let $\mathscr{C} = \{V_n : n \in \mathbb{N}\}$ be a local π-base at e. Put $\mathscr{F} = \{V_nV_n^{-1} : n \in \mathbb{N}\}$. Then $\mathscr{F}' = \{int(V_nV_n^{-1}) : n \in \mathbb{N}\}$ is a local base at e in τ^*.

Indeed, for each $n \in \mathbb{N}$ and fix a point $x \in V_n$, then $x^{-1}V_n$ is an open neighborhood at e in τ, and hence there exists an $U_\alpha \in \mathscr{B}$ such that $U_\alpha \subset x^{-1}V_n$. Thus $U_\alpha U_\alpha^{-1} \subset x^{-1}V_nV_n^{-1}x = V_nV_n^{-1}$, which implies that $V_nV_n^{-1}$ is a neighborhood of e in τ^*. On the other hand, fix $\alpha < \kappa$, there is $n \in \mathbb{N}$ such that $V_n \subset U_\alpha$. $V_nV_n^{-1} \subset U_\alpha U_\alpha^{-1}$. Therefore, $\mathscr{F}' = \{int(V_nV_n^{-1}) : n \in \mathbb{N}\}$ is a local base at e in τ^*.

Since first-countable topological group metrizable, we have (G, τ^*) is metrizable. Therefore, G is submetrizable. □

Since every submetrizable space is (hereditarily) Dieudonné complete, Corollary 3.9 give a partial answer to Question 1.6.

Theorem 3.11. *Let (G, τ) be a Hausdorff separable SIN first-countable paratopological group. There exists a continuous isomorphism of G onto a Tychonoff second-countable topological group.*

Proof. By the proof of Theorem 3.8, we know that (G, τ^*) is metrizable. Since G is separable and $\tau^* \subset \tau$, (G, τ^*) is separable, and hence (G, τ^*) is a second-countable topological group. Hence there exists a continuous isomorphism of G onto a Tychonoff second-countable topological group. □

By Theorem 3.5, we have the following corollary, which gives a partial answer to Question 1.7.

Corollary 3.12. *Let (G, τ) be a Hausdorff ω-narrow first-countable SIN paratopological group. There exists a continuous isomorphism of G onto a Tychonoff second-countable topological group.*

The following two theorems give another answers to Questions 1.4 and 1.6.

Theorem 3.13. *If (G, τ) is a Hausdorff saturated*[3] *first-countable paratopological group, then G is submetrizable.*

Proof. Suppose that $\{U_n : n \in \mathbb{N}\}$ is a countable local base at e. Let

$$\sigma = \{U \subset G : \text{There exists an } n \in \mathbb{N} \text{ such that } xU_nU_n^{-1} \subset U \text{ for each } x \in U\}.$$

Since G is saturated, it follows from [13, Theorem 3.2] that (G, σ) is a topological group. Obvious, (G, σ) is T_1 since (G, τ) is a Hausdorff, and hence (G, σ) is regular. Then (G, σ) is first-countable, and thus it is metrizable. Therefore, (G, τ) is submetrizable. □

Theorem 3.14. *If (G, τ) is a Hausdorff feebly compact*[4] *first-countable paratopological group, then G is submetrizable.*

Proof. Suppose that $\{U_n : n \in \mathbb{N}\}$ is a local base at e. Then the family $\{\mathrm{int}\overline{U_n} : n \in \mathbb{N}\}$ is a local base at e for a regular paratopological group topology σ on G. Obviously, (G, σ) is feebly compact. Since (G, σ) is first-countable, (G, σ) has a regular G_δ-diagonal [22]. Then (G, σ) is metrizable since a regular feebly compact space with a regular G_δ-diagonal is metrizable. Therefore, (G, τ) is submetrizable. □

Next, we gives an another answer to Question 1.2.

Theorem 3.15. *Let (G, τ) be a Hausdorff SIN paratopological group. If G is locally countable, then G is submetrizable.*

Proof. Since G is locally countable, there exists an open neighborhood U of e such that U is a countable set. Then UU^{-1} is also a countable set. Let $UU^{-1} \setminus \{e\} = \{x_n : n \in \mathbb{N}\}$. Since G is a paratopological group, we can find a family of countably many neighborhoods $\{V_n : n \in \omega\}$ of e satisfying the following conditions:
(i) $V_0 = U$;
(ii) For each $n \in \omega$, then $V_{n+1}^2 \subset V_n$;
(iii) For each $n \in \mathbb{N}$, then $x_n \notin V_nV_n^{-1}$ (This is possible since G is Hausdorff.)
(iv) For each $n \in \omega$, then $xV_{n+1}x^{-1} \subset V_n$ for each $x \in G$.

Since G is a SIN group, the topology σ generated by the neighborhood basis $\{V_n : n \in \mathbb{N}\}$ is a paratopological group. Clearly, (G, σ) is coarser than τ. Since $\bigcap_{n\in\mathbb{N}} V_nV_n^{-1} = \{e\}$ by (iii), σ is Hausdorff. Therefore, (G, σ) is a Hausdorff SIN first-countable paratopological group, and it follows from Theorem 3.8 that G is submetrizable. □

Corollary 3.16. *Let (G, τ) be a Hausdorff Abelian paratopological group. If G is locally countable, then G is submetrizable.*

---

[3] A paratopological group G is *saturated* if for any neighborhood U of e the set U^{-1} has nonempty interior in G.

[4] A space X is called *feebly compact* if each locally finite family of open subsets in X is finite.

4. Developable paratopological groups

Recall that a topological space is *symmetrizable* if its topology is generated by a symmetric, that is, by a distance function satisfying all the usual restrictions on a metric, except for the triangle inequality [1].

Now, we give a negative answer[5] to Question 1.8 by modifying [23, Example 2.1].

Example 4.1. There exists a separable, Moore paratopological group G such that G is not metrizable.

Proof. Let $G = \mathbb{R} \times \mathbb{Q}$ be the group with the usual addition. Then we define a topology on G by giving a local base at the neutral element $(0, 0)$ in G. For each $n \in \mathbb{N}$, let

$$U_n(0,0) = \{(0,0)\} \cup \{(x,y) : y \geq nx, y < \frac{1}{n}, y \in \mathbb{Q}, x \geq 0\}.$$

Let σ be the topology generated by the local base $\{U_n : n \in \mathbb{N}\}$ at the neutral element $(0, 0)$. It is easy to see that (G, σ) is a semitopological group. Now, we prove that is a paratopological group. Since G is Abelian, it only need to prove that for each $n \in \mathbb{N}$ there exists an $m \in \mathbb{N}$ such that $U_m^2 \subset U_n$. Indeed, fix an $n \in \mathbb{N}$, Then we have $U_{4n}^2 \subset U_n$. For two points $(x_i, y_i) \in U_{4n} (i = 1, 2)$, where $x_i \geq 0, y_i < \frac{1}{4n}, y_i \geq 4nx_i, y_i \in \mathbb{Q} (i = 1, 2)$. Let

$$(x, y) = (x_1, y_1) + (x_2, y_2) = (x_1 + x_2, y_1 + y_2).$$

Obviously, we have

$$x = x_1 + x_2 \geq 0, y = y_1 + y_2 < \frac{1}{4n} + \frac{1}{4n} = \frac{1}{2n} < \frac{1}{n},$$

and

$$y = y_1 + y_2 \geq 4nx_1 + 4nx_2 = 4nx \geq nx.$$

Then $(x, y) \in U_n$, and hence $U_{4n}^2 \subset U_n$. Moreover, it is easy to see that G is regular, separable and first-countable space.

For each $q \in \mathbb{Q}$, it is easy to see that the family $\{\{(x, q)\} : x \in \mathbb{R}\}$ is uncountable discrete and closed, hence G is a σ-space, and thus it is a β-space[6]. G is a Moore space by [23, Corollary 2.1]. Hence G is semi-metrizable [17]. Therefore G is symmetrizable since a space is semi-metrizable if only if it is first-countable and symmetrizable [17]. However, G is not metrizable since G is separable and contains a uncountable discrete closed subset. ☐

Question 4.2. *Is every quasi-developable paratopological (semitopological) group a β-space?*

Next, we give a partial answer to Question 4.2.

Lemma 4.3. [7, Lemma 1.2] *Suppose that G is a paratopological group and not a topological group. Then there exists an open neighborhood U of the neutral element e of G such that $U \cap U^{-1}$ is nowhere dense in G, that is, the interior of the closure of $U \cap U^{-1}$ is empty.*

---

[5] Li, Mou and Wang [19] also obtained a non-metrizable Moore paratopological group.

[6] Let (X, τ) be a topological space. A function $g : \omega \times X \to \tau$ satisfies that $x \in g(n, x)$ for each $x \in X, n \in \omega$. A space X is a β-space [17] if there is a function $g : \omega \times X \to \tau$ such that if $x \in g(n, x_n)$ for each $n \in \omega$, then the sequence $\{x_n\}$ has a cluster point in X.

Theorem 4.4. *A regular Baire*[7] *quasi-developable paratopological group G is a metrizable topological group.*

Proof. Claim: Let U be an arbitrary open neighborhood of e. Then $\overline{U^{-1}}$ is also a neighborhood of e.

Suppose that $\{\mathscr{U}_n : n \in \mathbb{N}\}$ is a family of open subsets of G such that for each $x \in G$ and $x \in V$ with V open in G, there exists an $n \in \mathbb{N}$ such that $x \in \mathrm{st}(x, \mathscr{U}_n) \subset V$. For each $\mathbb{N}$, put

$$A_n = \{x \in G : \mathrm{st}(x, \mathscr{U}_n) \subset x \cdot U\}.$$

It is easy to see that $G = \bigcup\{A_n : n \in \mathbb{N}\}$. Since G is Baire, there exists $n_0 \in \mathbb{N}$ such that $\mathrm{int}\overline{A_{n_0}} \neq \emptyset$. Therefore, there exist a point $x_0 \in G$ and $n_1 \in \mathbb{N}$ such that $\mathrm{st}(x_0, \mathscr{U}_{n_1}) \subset \overline{A_{n_0}}$. Let $\mathscr{V} = \{U_1 \cap U_2 : U_1 \in \mathscr{U}_{n_0}, U_2 \in \mathscr{U}_{n_1}\}$. Put $W = \mathrm{st}(x_0, \mathscr{V})$. For each $W \cap A_{n_0}$, it is easy to see

$$x_0 \in \mathrm{st}(y, \mathscr{V}) \subset \mathrm{st}(y, \mathscr{U}_{n_0}) \subset y \cdot U,$$

hence $y^{-1}x_0 \in U$, so $x_0^{-1}y \in U^{-1}$, hence $x_0^{-1} \cdot (W \cap A_{n_0}) \subset U^{-1}$. Moreover, since $W \subset \overline{A_{n_0}}$, then $W \subset \overline{W \cap A_{n_0}}$. Therefore, we have

$$e \in x_0^{-1}W \subset x_0^{-1} \cdot \overline{W \cap A_{n_0}} \subset \overline{x_0^{-1} \cdot (W \cap A_{n_0})} \subset \overline{U^{-1}}.$$

Since $x_0^{-1}W$ is an open neighborhood of e, the set $\overline{U^{-1}}$ is also a neighborhood of e.

$\overline{U^{-1}} \cap U \subset \overline{U \cap U^{-1}}$, by Claim, $\overline{U \cap U^{-1}}$ contains a neighborhood of e, then $U \cap U^{-1}$ is not nowhere dense. By Lemma 4.3, G is a topological group. Therefore G is metrizable since first-countable topological groups are metrizable. □

Finally, we pose some questions about developable paratopological groups.

Question 4.5. *Is each regular Baire quasi-developable semitopological group G is a paratopological group?*

If G is developable, then the answer is affirmative. Indeed, it was proved that each Baire Moore semitopological group G is a metrizable topological group [14].

Question 4.6. *Is every developable or Moore paratopological group submetrizable?*

Question 4.7. *Is every normal Moore paratopological group submetrizable?*

Question 4.8. *Is every paratopological group with a base of countable order*[8] *developable?*

---

[7] Recall that a space is *Baire* if the intersection of a sequence of open and dense subsets is dense.

[8] A space X is said to have a *base of countable order*(BCO) [17] if there is a sequence $\{\mathcal{B}_n\}$ of base for X such that whenever $x \in b_n \in \mathcal{B}_n$ and (b_n) is decreasing (by set inclusion), then $\{b_n : n \in \mathbb{N}\}$ is a base at x.

5. Fréchet-Urysohn paratopological groups

First, we need the following Lemma.

Lemma 5.1. [8, Theorem 4.7.5] *Every weakly first-countable Hausdorff paratopological group is first-countable.*

Arhangel'skii proved that if a topological group G is an image of a separable metrizable space under a pseudo-open[9] map, then G is metrizable [4]. We have the following.

Theorem 5.2. *Let G be a uncountable paratopological group. Suppose that G is a pseudoopen image of a separable metric space, then G is a separable and metrizable.*

Proof. We introduce a new product operation in the topological space G by the formula: $a \times b = ba$, for $a, b \in G$ and denote the space with this operation by H. Put $T = \{(g, g^{-1}) \in G \times H, g \in G\}$. $|T| > \omega$ since G is uncountable. By [7, Proposition 2.9], H is a paratopological group and T is closed in the space $G \times H$ and is a topological group.

Since G is a pseusdo-open image of a separable metric space, then the space G is a Fréchet-Urysohn space with a countable k-network. $G \times H$ has a countable network. Hence T has a countable network. By the proof of [17, Theorem 4.9], T is a one-to-one continuous image of a separable metric space M. Let D be a countable dense subset of M, there is a sequence $L \subset D$ converging to some point in $M \setminus D$ since M is uncountable. Therefore, there is a non-trivial sequence $\{(g_n, g_n^{-1}) : n \in \mathbb{N}\}$ converging to (e, e) (note that T is homogeneous), and hence there exists a sequence $C_0 = \{g_n : n \in \mathbb{N}\} \subset G$ converging to e and its inverse $C_1 = \{g_n^{-1} : n \in \mathbb{N}\}$ also converges to e. G contains no closed copy of S_2 since G is Fréchet-Urysohn. By [22, Theorem 2.4], G contains no closed copy of S_ω. Since G is a sequential space with a point-countable k-network and contains no closed copy of S_ω, then G is weakly first-countable [20], and hence G is first-countable by Lemma 5.1. Therefore G is separable and metrizable [17, Theorem 11.4(ii)]. □

A quotient image of a topological sum of countably many compact spaces is called a k_ω-space. Every countable k_ω-space is a sequential $\aleph_0$-space, and a product of two k_ω-spaces is itself a k_ω-space, see [26].

Theorem 5.3. *Let G be a regular countable, locally k_ω, paratopological group. Then G is a discrete topological group or contains a closed copy of S_ω.*

Proof. Suppose that G is an α_4-space. Then since G is locally k_ω, G is sequential and $\aleph_0$, and thus G is weakly first-countable [21]. Then G is first-countable by Lemma 5.1, and hence G is a separable metrizable space since G is countable. If G is not discrete, G has no isolated points. Then G is homeomorphic to the rational number set $\mathbb{Q}$ since a separable metrizable space is homeomorphic to the rational number set $\mathbb{Q}$ provided that it is infinite, countable and without any isolated points [18]. However, $\mathbb{Q}$ is not a locally k_ω-space, which is a contradiction. Then G is discrete, and G is a topological group.

If G is not an α_4-space, and thus G contains a copy of S_ω. Since every point of G is a G_δ-set, it follows from [20, Corollary 3.4] that G contains a closed copy of S_ω. □

---

[9] A map $f : X \to Y$ is pseudo-open if for each $y \in Y$ and every open set U containing $f^{-1}(y) \subset U$ one has $y \in int(f(U))$.

By Theorem 5.3, it is easy to obtain the following corollary.

Corollary 5.4. *Let G be a regular, countable, non-discrete, Fréchet-Urysohn*[10] *paratopological group. If G is k_ω, then G contains a closed copy of S_ω and no closed copy of S_2.*

Since the closed image of a locally compact, separable metric space is a Fréchet-Urysohn and k_ω-space, we have the following corollary.

Corollary 5.5. *Let G be a countable non-discrete paratopological group. If G is a closed image of a locally compact, separable metric space, then G contains a closed copy of S_ω, and hence G is not metrizable.*

The condition "locally k_ω-space" is essential in Theorem 5.3, and we can not omit it.

Example 5.6. There exists a regular non-discrete countable second-countable paratopological group G such that G contains no closed copy of S_ω and G is not a topological group.

Proof. Let $G = \mathbb{Q}$ be the rational number and endow with the subspace topology of the Sorgenfrey line. Then $\mathbb{Q}$ is a second-countable paratopological group and non-discrete. Obviously, G contains no closed copy of S_ω and G is not a topological group. □

6. NON-H-CLOSED PARATOPOLOGICAL GROUPS

A paratopological group is H-closed if it is closed in every Hausdorff paratopological group containing it as a subgroup.

Let U be a neighborhood of e in a paratopological group G. We say that a subset $A \subset G$ is U-unbounded if $A \nsubseteq KU$ for every finite subset $K \subset G$.

Now, we give a negative answer to Question 1.10.

Lemma 6.1. [28] *Let G be an abelian paratopological group of the infinite exponent. If there exists a neighborhood U of the neutral element such that a group nG is UU^{-1}-unbounded for every $n \in \mathbb{N}$, then the paratopological group G is not H-closed.*

Theorem 6.2. *The Sorgenfrey line $(\mathbb{R}, \tau)$ is not H-closed.*

Proof. Obvious, $\mathbb{R}$ is an abelian paratopological group of the infinite exponent. Let $U = [0, 1)$. Then $UU^{-1} = (-1, 1)$ For each $n \in \mathbb{N}$, it is easy to see that $n\mathbb{R} \nsubseteq K(-1, 1)$ for every finite subset K of $\mathbb{R}$. Indeed, for each finite subset K of $\mathbb{R}$, since K is finite set, there exists an $n \in \mathbb{N}$ such that $|x| \leq n$ for each $x \in K$, and then $K \subset (-1, 1) \subset (-n, n)$. Therefore, we have $n\mathbb{R} \nsubseteq K(-1, 1)$. By Lemma 6.1, $\mathbb{R}$ is not H-closed. □

However, we have the following theorem.

Theorem 6.3. *Let $(\mathbb{R}, \tau)$ be the Sorgenfrey line. Then the quotient group $(\mathbb{R}/\mathbb{Z}, \xi)$ is H-closed, where ξ is the quotient topology.*

---

[10] A space X is said to be *Fréchet-Urysohn* if, for each $x \in \overline{A} \subset X$, there exists a sequence $\{x_n\}$ such that $\{x_n\}$ converges to x and $\{x_n : n \in \mathbb{N}\} \subset A$.

Proof. Let $(\mathbb{R}/\mathbb{Z}, \sigma)$ be the finest group topology such that $\sigma \subset \xi$. Then $(\mathbb{R}/\mathbb{Z}, \sigma)$ is compact. Therefore, $(\mathbb{R}/\mathbb{Z}, \sigma)$ is H-closed. Hence $(\mathbb{R}/\mathbb{Z}, \xi)$ is H-closed by [28, Proposition 10]. □

Let G be an abelian non-periodic paratopological group. We say that G is *strongly unbounded* if there exists non-periodic element x_0 and open neighborhood U of e such that $\langle x_0\rangle \cap UU^{-1} = \{e\}$ and $\langle x_0\rangle$ is closed in G, where $\langle x_0\rangle$ is a subgroup generated by x_0. Obviously, every strongly unbounded paratopological group is not H-closed.

Next, we discuss some non-H-closed paratopological groups.

Given any elements $a_0, a_1, \cdots, a_n$ of an abelian group G put

$$X(a_0, a_1, \cdots, a_n) = \{a_0^{x_0} a_1^{x_1} \cdots a_n^{x_n} : 0 \leq x_i \leq n, 0 \leq i \leq n\}.$$

Theorem 6.4. *Let (G, τ) be a Hausdorff strongly unbounded paratopological group. Then there exists a Tychonoff paratopological group topology γ on $G \times \mathbb{Z}$ satisfies the following conditions:*

1. *There exists a Hausdorff paratopological group topology $\sigma \subset \gamma$ on $G \times \mathbb{Z}$ such that $\sigma|_{G\times\{0\}} = \tau$ and $(G \times \mathbb{Z}, \sigma)$ contains closed copies of S_2 and S_ω;*
2. *$(G \times \mathbb{Z}, \gamma)$ is a strongly zero-dimensional, paracompact σ-space;*
3. *The remainder $b(G \times \mathbb{Z}, \gamma) \setminus (G \times \mathbb{Z}, \gamma)$ of every Hausdorff compactification $b(G \times \mathbb{Z}, \gamma)$ is pseudocompact.*

Proof. Since G is strongly unbounded paratopological group, there exist a non-periodic element x_0 of G and open neighborhood U of e such that $\langle x_0\rangle \cap UU^{-1} = \{e\}$ and $\langle x_0\rangle$ is closed in G. Obvious, $\langle x_0\rangle$ is an abelian group. Then the mapping $f : (nx_0, m) \mapsto (n, m)$ is is naturally isomorphic from $\langle x_0\rangle \times \mathbb{Z}$ onto a group $\mathbb{Z} \times \mathbb{Z}$. We may assume that $\mathbb{Z} \times \mathbb{Z} \subset G \times \mathbb{Z}$. Now we define a zero dimensional paratopological group topology on $G \times \mathbb{Z}$. Obvious, we can define a positively natural number sequence $\{a_n\}$ satisfies the following conditions:
(1) $a_n > n$;
(2) $a_n > 2a$ for each $a \in X(a_1, \cdots, a_{n-1})$.
Define a base $\mathscr{B}_\gamma$ at the neutral element of paratopological group topology γ on the group $G \times \mathbb{Z}$ as follows. Put $A_n^+ = \{(e, 0)\} \cup \{(a_k, 1) : k > n\}$. For every strictly increasing sequence $\{n_k\}$ put $A[n_k] = \bigcup_{l\in\mathbb{N}} A_{n_1}^+ \cdots A_{n_l}^+$. Put $\mathscr{B}_\gamma = \{A[n_k]\}$. Then γ is a zero dimensional paratopological group topology on $G \times \mathbb{Z}$, see [28, Lemma 3].

(1) By the proof of [28, Lemma 3], we can define a topology σ on $G \times \mathbb{Z}$ such that $\sigma|_{G\times\{0\}} = \tau$, $\sigma \subset \gamma$ and $\sigma|_{\mathbb{Z}\times\mathbb{Z}} = \gamma|_{\mathbb{Z}\times\mathbb{Z}}$. Let $\gamma|_{\mathbb{Z}\times\mathbb{Z}} = \xi$. Since $(\mathbb{Z} \times \mathbb{Z}, \xi)$ is zero dimensional and countable, the space $(\mathbb{Z} \times \mathbb{Z}, \xi)$ is Tychonoff.

Claim 1: $(\mathbb{Z} \times \mathbb{Z}, \xi)$ is a closed in $(G \times \mathbb{Z}, \sigma)$.

Since $\langle 1\rangle \cap UU^{-1} = \{e\}$ and $\langle 1\rangle$ is closed in G, it is easy to see that $(\mathbb{Z} \times \mathbb{Z}, \xi)$ is closed in $(G \times \mathbb{Z}, \sigma)$.

Claim 2: $(\mathbb{Z} \times \mathbb{Z}, \xi)$ contains a closed copy of S_ω.

For each $n \in \mathbb{N}$, let $\beta_n = \{(na_{n+k}, n)\}_{k\in\mathbb{N}}$. Obvious, each β_n converges to $(0, 0)$ as $k \to \infty$.

Let $X = \{(na_{n+k}, n) : k, n \in \mathbb{N}\} \cup \{(0, 0)\}$. It is easy to see that X is a closed copy of S_ω.

Claim 3: $(\mathbb{Z} \times \mathbb{Z}, \xi)$ contains a closed copy of S_2.

Let $\alpha_0 = \{(a_k, 1)\}_{k\in\mathbb{N}}$. For each $n \in \mathbb{N}$, let $\alpha_n = \{(a_n + (n-1)a_k, n)\}_{k\in\mathbb{N}}$. Obvious, α_0 converges to $(0,0)$ as $k \to \infty$ and each α_n converges to $(a_n, 1)$ as $k \to \infty$.

Let $X = \{(a_n + (n-1)a_k, n) : k, n \in \mathbb{N}\} \cup \alpha_0 \cup \{(0,0)\}$. It is easy to see that X is a closed copy of S_2.

(2) Let $i : \mathbb{Z} \to \mathbb{Z}$ be the identity map. Since $\mathbb{Q}$ is a divisible group, the map i can be extended to a homomorphism $\phi : G \to \mathbb{Q}$. Put $|x| = |\phi(x)|$ for every element $x \in G$. Then $|\cdot|$ is a seminorm on the group G such that for all $x, y \in G$ holds $|x+y| \leq |x| + |y|$. For each $n \in \mathbb{N}$, let $\mathscr{B}_n = \{\{(x,y)\} : \phi(x) \leq n, y \in \mathbb{Z}\}$. Then it is easy to see that each $\mathscr{B}_n$ is a discrete family of closed subsets. Since it follows from Theorem 3.15 that the space $(G \times \mathbb{Z}, \gamma)$ is submetrizable, there exists a metrizale topology $\mathscr{F}$ such that $\mathscr{F} \subset \gamma$. Obvious, $\bigcup_{n\in\mathbb{N}} \mathscr{B}_n$ is a network for $(G\times\mathbb{Z}, \mathscr{F})$, and it follows from [8, Theorem 7.6.6] that $(G\times\mathbb{Z}, \gamma)$ is a paracompact σ-space. Moreover, each subspace $F_n = \bigcup \mathscr{B}_n$ is strongly zero-dimensional and $G \times \mathbb{Z} = \bigcup_{n\in\mathbb{N}} F_n$, and hence $(G \times \mathbb{Z}, \gamma)$ is strongly zero-dimensional by [15, Theorem 2.2.7].

(3) **Claim 4**: $(G \times \mathbb{Z}, \gamma)$ has no strong π-base[11] at any compact subset of $G \times \mathbb{Z}$.

Indeed, suppose there exists a compact subset $K \subset G \times \mathbb{Z}$ such that K has a strong π-base φ. It is easy to see that the set $\{n : K \cap (G \times \{n\}) \neq \emptyset, n \in \mathbb{Z}\}$ is finite. Moreover, without loss of generalization, we may assume that strong π-base φ at K is countable and each element of φ is an open neighborhood of some point in $G\times\mathbb{Z}$. Let $A = \{(b_i, n(i)) : i \in \mathbb{N}, n(i) \in \mathbb{Z}\}$, and let $\varphi = \{(b_i, n(i)) + A[n_k^i] : i \in \mathbb{N}\}$.

Case 1: The set $\{n(i) : i \in \mathbb{N}\}$ is infinite.

Then it is easy to see that A contains a discrete closed subset B such that $B \cap K = \emptyset$. Let U be an open neighborhood at K. Then $U \setminus B$ is also an open neighborhood at K, and each $(b_i, n(i)) + A[n_k^i] \nsubseteq U \setminus B$, which is a contradiction.

Case 2: The set $\{n(i) : i \in \mathbb{N}\}$ is finite.

Let $n_0 = \max(\{n(i) : i \in \mathbb{N}\} \bigcup \{n : K \cap (G\times\{n\}) \neq \emptyset, n \in \mathbb{Z}\} \bigcup \{0\}) + 1$. For each $i + n_0 \in \mathbb{N}$, we can choose a $(c_i, i+n_0) \in (b_i, n(i)) + A[n_k^i]$. Then $\{(c_i, i+n_0) : i \in \mathbb{N}\}$ is closed and discrete subset in $(G \times \mathbb{Z}, \gamma)$ and $\{(c_i, i+n_0) : i \in \mathbb{N}\} \cap K = \emptyset$. Let U be an open neighborhood at K. Then $U \setminus \{(c_i, i+n_0) : i \in \mathbb{N}\}$ is also an open neighborhood at K, and each $(b_i, n(i)) + A[n_k^i] \nsubseteq U \setminus \{(c_i, i+n_0) : i \in \mathbb{N}\}$, which is a contradiction.

It follows from Claim 4 and [6, Corollary 4.3] that $b(G \times \mathbb{Z}) \setminus (G \times \mathbb{Z})$ is pseudocompact. $\square$

Remark (1) Sorgenfrey line $(\mathbb{R}, \tau)$ is a Hausdorff strongly unbounded paratopological group, and hence we can define a Hausdorff paratopological group topology σ on $G \times \mathbb{Z}$ such that $\sigma|_{G\times\{0\}} = \tau$ and $(G \times \mathbb{Z}, \sigma)$ contains closed copies of S_2 and S_ω;

(2) Let $\mathbb{Q}$ be the rationals with the subspace topology of usual topology $\mathbb{R}$. Then $\mathbb{Q}$ is a topological group. It easily check that $\mathbb{Q}$ is a strongly zero-dimensional, nowhere locally compact, paracompact σ-space. In additional, $\mathbb{Q}$ has a strong π-base at each point since it is first-countable. It follows from [3, Lemma 2.1] that the remainder of any Hausdorff compactification of $\mathbb{Q}$ is not pseudocompact.

A paratopological group G is said to have the property $(\star\star)$, if there exists a sequence $\{x_n : n \in \mathbb{N}\}$ of G such that $x_n \to e$ and $x_n^{-1} \to e$.

---

[11] A *strong π-base* of a space X at a subset F of X is an infinite family γ of non-empty open subsets of X such that every open neighborhood of F contains all but finitely many elements of γ. Clearly, a strong π-base can be always assumed to be countable.

In [22], C. Liu proved the following theorem.

Theorem 6.5. [22] *Let G be a paratopological group having the property (**). Then G has a (closed) copy of S_2 if it has a (closed) copy of S_ω.*

It is natural to ask the following.

Question 6.6. [22] *Can we omit the property (**) in Theorem 6.5.*

By Lemma 6.1, the space G in Theorem 6.4 is not H-closed. Moreover, it is easy to see that the paratopological group topology on $\mathbb{Z} \times \mathbb{Z}$ in Theorem 6.4 does not have the the property (**). Then we have the following questions.

Question 6.7. *Let (G, τ) be a H-closed paratopological group. Does there exist a Hausdorff paratopological group topology σ on $G \times \mathbb{Z}$ such that $\sigma|_{G\times\{0\}} = \tau$ and $(G \times \mathbb{Z}, \sigma)$ contains closed copies of S_2 and S_ω?*

Question 6.8. *Let G be a not H-closed paratopological group. Is it true that G has a (closed) copy of S_2 if it has a (closed) copy of S_ω?*

Question 6.9. *Let G be a H-closed paratopological group. Is it true that G has a (closed) copy of S_2 if it has a (closed) copy of S_ω?*

Acknowledgements. We wish to thank the reviewers for the detailed list of corrections, suggestions to the paper, and all her/his efforts in order to improve the paper.

References

[1] A.V. Arhangel'skiĭ, Mappings and spaces, *Russian Math. Surveys*, 1966, **21**, 115–162.

[2] A.V. Arhangel'skiĭ, *Topological invariants in algebraic enviroment*, Recent progress in General Topology II, ed. by Hušek and van Mill, ELSEVIER, (2002), 1–58.

[3] A. Arhangel'skiĭ, *Two types of remainders of topological groups*, Comment. Math. Univ. Carolin., 49(2008), 119–126.

[4] A.V. Arhangel'skiĭ, *Some classes of quotient and pseudoopen mappings and some new cardinal invariants of tightness type*, International Conference on Topology and its Applications, Islamabad, Pakistan, July 4–10, 2011.

[5] A.V. Arhangel'skiĭ, D.K. Burke, *Spaces with a regular G_δ-diagonal*, Topology Appl., **153**(2006), 1917–1929.

[6] A.V. Arhangel'skiĭ, M.M. Choban, *Remainders of rectifiable spaces*, Topology Appl., **157**(2010), 789–799.

[7] A.V. Arhangel'skiĭ, E.A. Reznichenko, *Paratopological and semitopological groups versus topological groups*, Topology Appl., **151**(2005), 107–119.

[8] A.V. Arhangel'skiĭ, M. Tkachenko, *Topological Groups and Related Structures*, Atlantis Press and World Sci., 2008.

[9] D. Basile, A. Bella, G.J. Ridderbos, *Weak extent, submetrizability and diagonal degrees*, arXiv:1112.0883v1[math.GN].

[10] H.R. Bennett, *On quasi-developable spaces*, General Topology Appl., **1**(1971), 253–262.

[11] H.R. Bennett, D. Lutzer, *Ordered spaces with special bases*, Fundamenta Mathematicae, **158**(1998), 289–299.

[12] H.R. Bennett, D. Lutzer, *Recent developments in the topology of ordered spaces*, Recent progress in General Topology II, ed. by Hušek and van Mill, ELSEVIER, (2002), 85-114.

[13] T. Banakh, I. Guran, O. Ravsky, *Characterizing meager paratopological groups*, Applied General Topology, **12(1)**(2011), 27–33.

[14] J. Cao, R. Drozdowski, Z. Piotrowski, *Weak continuity properties of topological groups*, Czech. Math. J., **60(135)**(2010), 133–148.

[15] R. Engelking, *Dimension theory*, PWN, Polish Scientific Publ., Warszawa, 1978.

[16] R. Engelking, *General Topology* (revised and completed edition), Heldermann Verlag, Berlin, 1989.

[17] G. Gruenhage, *Generalized metric spaces*, In: K. Kunen, J. E. Vaughan(Eds.), Handbook of Set-Theoretic Topology, Elsevier Science Publishers B.V., Amsterdam, 1984, 423–501.
[18] K.P. Hart, J. Nagata, J.E. Vaughan, *Encyclopedia of General topology*, Elsevier Science Publishers B.V., Amsterdam, 2004, 337–340.
[19] P.Y. Li, L. Mou, S.Z. Wang, *Notes on questions about spaces with algebraic strucures*, to appear in Topology Appl.
[20] S. Lin, A note on the Arens' space and sequential fan, *Topology Appl.*, **81**(1997), 185–196.
[21] S. Lin, Point-countable Covers and Sequence-covering Mappings(in Chinese): Science Press, Beijing, 2002.
[22] C. Liu, *A note on paratopological groups*, Comment.Math.Univ.Carolin., **47**(2006), 633–640.
[23] C. Liu, S. Lin, *Generalized metric spaces with algebraic structures*, *Topology Appl.*, **157**(2010), 1966–1974.
[24] C. Liu, *Metrizability of paratopological (semitopological) groups*, *Topology Appl.*, **159**(2012), 1415–1420.
[25] D. Lutzer, *On generalized ordered spaces*, Dissertationes Math., 89 (1971).
[26] E. Michael, *A quintuple quotient quest*, *Gen. Topology Appl.*, **2**(1972), 91–138.
[27] N.M. Pyrch, A.V. Ravsky, *On free paratopological groups*, Matematychni Studii, **25**(2006), 115–125.
[28] O. Ravsky, *On H-closed paratopological groups*, arXiv:1003.5377v1.
[29] O. Ravsky, *Paratopological groups I*, Mathematychni Studii, **16**(2001), 37-48.
[30] J.M. Worrell, H.H. Wicke, *Characterizations of developable topological spaces*, *Canad. J. Math.*, **17**(1965), 820–830.

(Fucai Lin): Department of Mathematics and Information Science, Zhangzhou Normal University, Zhangzhou 363000, P. R. China
E-mail address: `linfucai2008@yahoo.com.cn`

(Chuan Liu): Department of Mathematics, Ohio University Zanesville Campus, Zanesville, OH 43701, USA
E-mail address: `liuc1@ohio.edu`